

Intended for *Phys. Rev. D*

Authors: Y. P. Guo, C. Z. Yuan, C. X. Yu

Committee: M. Maggiora (Chair), F. Liu, X. T. Huang

Search for hadronic transition $\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-$ and observation of $\chi_{cJ} \rightarrow K \bar{K} \pi \pi$

M. Ablikim¹, M. N. Achasov⁵, D. J. Ambrose³⁹, F. F. An¹, Q. An⁴⁰, Z. H. An¹, J. Z. Bai¹,
Y. Ban²⁷, J. Becker², M. Bertani^{18A}, J. M. Bian³⁸, E. Boger^{20,a}, O. Bondarenko²¹, I. Boyko²⁰,
R. A. Briere³, V. Bytev²⁰, X. Cai¹, O. Cakir^{35A}, A. Calcaterra^{18A}, G. F. Cao¹, S. A. Cetin^{35B},
J. F. Chang¹, G. Chelkov^{20,a}, G. Chen¹, H. S. Chen¹, J. C. Chen¹, M. L. Chen¹, S. J. Chen²⁵,
Y. B. Chen¹, H. P. Cheng¹⁴, Y. P. Chu¹, D. Cronin-Hennessy³⁸, H. L. Dai¹, J. P. Dai¹,
D. Dedovich²⁰, Z. Y. Deng¹, A. Denig¹⁹, I. Denysenko^{20,b}, M. Destefanis^{43A,43C}, W. M. Ding²⁹,
Y. Ding²³, L. Y. Dong¹, M. Y. Dong¹, S. X. Du⁴⁶, J. Fang¹, S. S. Fang¹, L. Fava^{43B,43C},
F. Feldbauer², C. Q. Feng⁴⁰, R. B. Ferrolri^{18A}, C. D. Fu¹, J. L. Fu²⁵, Y. Gao³⁴, C. Geng⁴⁰,
K. Goetzen⁷, W. X. Gong¹, W. Gradl¹⁹, M. Greco^{43A,43C}, M. H. Gu¹, Y. T. Gu⁹, Y. H. Guan⁶,
A. Q. Guo²⁶, L. B. Guo²⁴, Y. P. Guo²⁶, Y. L. Han¹, F. A. Harris³⁷, K. L. He¹, M. He¹,
Z. Y. He²⁶, T. Held², Y. K. Heng¹, Z. L. Hou¹, H. M. Hu¹, J. F. Hu⁶, T. Hu¹, G. M. Huang¹⁵,
J. S. Huang¹², X. T. Huang²⁹, Y. P. Huang¹, T. Hussain⁴², C. S. Ji⁴⁰, Q. Ji¹, X. B. Ji¹,
X. L. Ji¹, L. L. Jiang¹, X. S. Jiang¹, J. B. Jiao²⁹, Z. Jiao¹⁴, D. P. Jin¹, S. Jin¹, F. F. Jing³⁴,
N. Kalantar-Nayestanaki²¹, M. Kavatsyuk²¹, W. Kuehn³⁶, W. Lai¹, J. S. Lange³⁶, C. H. Li¹,
Cheng Li⁴⁰, Cui Li⁴⁰, D. M. Li⁴⁶, F. Li¹, G. Li¹, H. B. Li¹, J. C. Li¹, K. Li¹⁰, Lei Li¹, Q. J. Li¹,
S. L. Li¹, W. D. Li¹, W. G. Li¹, X. L. Li²⁹, X. N. Li¹, X. Q. Li²⁶, X. R. Li²⁸, Z. B. Li³³,
H. Liang⁴⁰, Y. F. Liang³¹, Y. T. Liang³⁶, G. R. Liao³⁴, X. T. Liao¹, B. J. Liu¹, C. L. Liu³,
C. X. Liu¹, C. Y. Liu¹, F. H. Liu³⁰, Fang Liu¹, Feng Liu¹⁵, H. Liu¹, H. B. Liu⁶, H. H. Liu¹³,
H. M. Liu¹, H. W. Liu¹, J. P. Liu⁴⁴, K. Y. Liu²³, Kai Liu⁶, P. L. Liu²⁹, Q. Liu⁶, S. B. Liu⁴⁰,
X. Liu²², X. H. Liu¹, Y. B. Liu²⁶, Z. A. Liu¹, Zhiqiang Liu¹, Zhiqing Liu¹, H. Loehner²¹,
G. R. Lu¹², H. J. Lu¹⁴, J. G. Lu¹, Q. W. Lu³⁰, X. R. Lu⁶, Y. P. Lu¹, C. L. Luo²⁴, M. X. Luo⁴⁵,
T. Luo³⁷, X. L. Luo¹, M. Lv¹, C. L. Ma⁶, F. C. Ma²³, H. L. Ma¹, Q. M. Ma¹, S. Ma¹, T. Ma¹,
X. Y. Ma¹, Y. Ma¹¹, F. E. Maas¹¹, M. Maggiora^{43A,43C}, Q. A. Malik⁴², Y. J. Mao²⁷, Z. P. Mao¹,
J. G. Messchendorp²¹, J. Min¹, T. J. Min¹, R. E. Mitchell¹⁷, X. H. Mo¹, C. Morales
Morales¹¹, C. Motzko², N. Yu. Muchnoi⁵, H. Muramatsu³⁹, Y. Nefedov²⁰, C. Nicholson⁶,
I. B. Nikolaev⁵, Z. Ning¹, S. L. Olsen²⁸, Q. Ouyang¹, S. Pacetti^{18B}, J. W. Park²⁸, M. Pelizaeus³⁷,
H. P. Peng⁴⁰, K. Peters⁷, J. L. Ping²⁴, R. G. Ping¹, R. Poling³⁸, E. Prencipe¹⁹, M. Qi²⁵,
S. Qian¹, C. F. Qiao⁶, X. S. Qin¹, Y. Qin²⁷, Z. H. Qin¹, J. F. Qiu¹, K. H. Rashid⁴², G. Rong¹,
X. D. Ruan⁹, A. Sarantsev^{20,c}, B. D. Schaefer¹⁷, J. Schulze², M. Shao⁴⁰, C. P. Shen^{37,d},
X. Y. Shen¹, H. Y. Sheng¹, M. R. Shepherd¹⁷, X. Y. Song¹, S. Spataro^{43A,43C}, B. Spruck³⁶,
D. H. Sun¹, G. X. Sun¹, J. F. Sun¹², S. S. Sun¹, Y. J. Sun⁴⁰, Y. Z. Sun¹, Z. J. Sun¹, Z. T. Sun⁴⁰,
C. J. Tang³¹, X. Tang¹, I. Tapan^{35C}, E. H. Thorndike³⁹, D. Toth³⁸, M. Ullrich³⁶, G. S. Varner³⁷,
B. Wang⁹, B. Q. Wang²⁷, K. Wang¹, L. L. Wang⁴, L. S. Wang¹, M. Wang²⁹, P. Wang¹,
P. L. Wang¹, Q. Wang¹, Q. J. Wang¹, S. G. Wang²⁷, X. L. Wang⁴⁰, Y. D. Wang⁴⁰, Y. F. Wang¹,
Y. Q. Wang²⁹, Z. Wang¹, Z. G. Wang¹, Z. Y. Wang¹, D. H. Wei⁸, P. Weidenkaff¹⁹,
Q. G. Wen⁴⁰, S. P. Wen¹, M. Werner³⁶, U. Wiedner², L. H. Wu¹, N. Wu¹, S. X. Wu⁴⁰,
W. Wu²⁶, Z. Wu¹, L. G. Xia³⁴, Z. J. Xiao²⁴, Y. G. Xie¹, Q. L. Xiu¹, G. F. Xu¹, G. M. Xu²⁷,
H. Xu¹, Q. J. Xu¹⁰, X. P. Xu³², Z. R. Xu⁴⁰, F. Xue¹⁵, Z. Xue¹, L. Yan⁴⁰, W. B. Yan⁴⁰,
Y. H. Yan¹⁶, H. X. Yang¹, Y. Yang¹⁵, Y. X. Yang⁸, H. Ye¹, M. Ye¹, M. H. Ye⁴, B. X. Yu¹,

C. X. Yu²⁶, J. S. Yu²², S. P. Yu²⁹, C. Z. Yuan¹, Y. Yuan¹, A. A. Zafar⁴², A. Zallo^{18A}, Y. Zeng¹⁶,
 B. X. Zhang¹, B. Y. Zhang¹, C. C. Zhang¹, D. H. Zhang¹, H. H. Zhang³³, H. Y. Zhang¹,
 J. Q. Zhang¹, J. W. Zhang¹, J. Y. Zhang¹, J. Z. Zhang¹, S. H. Zhang¹, X. J. Zhang¹,
 X. Y. Zhang²⁹, Y. Zhang¹, Y. H. Zhang¹, Y. S. Zhang⁹, Z. P. Zhang⁴⁰, Z. Y. Zhang⁴⁴, G. Zhao¹,
 H. S. Zhao¹, J. W. Zhao¹, K. X. Zhao²⁴, Lei Zhao⁴⁰, Ling Zhao¹, M. G. Zhao²⁶, Q. Zhao¹,
 S. J. Zhao⁴⁶, T. C. Zhao¹, X. H. Zhao²⁵, Y. B. Zhao¹, Z. G. Zhao⁴⁰, A. Zhemchugov^{20,a},
 B. Zheng⁴¹, J. P. Zheng¹, Y. H. Zheng⁶, B. Zhong¹, J. Zhong², L. Zhou¹, X. K. Zhou⁶,
 X. R. Zhou⁴⁰, C. Zhu¹, K. Zhu¹, K. J. Zhu¹, S. H. Zhu¹, X. L. Zhu³⁴, X. W. Zhu¹,
 Y. C. Zhu⁴⁰, Y. M. Zhu²⁶, Y. S. Zhu¹, Z. A. Zhu¹, J. Zhuang¹, B. S. Zou¹, J. H. Zou¹

(BESIII Collaboration)

¹ *Institute of High Energy Physics, Beijing 100049, P. R. China*

² *Bochum Ruhr-University, 44780 Bochum, Germany*

³ *Carnegie Mellon University, Pittsburgh, PA 15213, USA*

⁴ *China Center of Advanced Science and Technology, Beijing 100190, P. R. China*

⁵ *G.I. Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia*

⁶ *Graduate University of Chinese Academy of Sciences, Beijing 100049, P. R. China*

⁷ *GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany*

⁸ *Guangxi Normal University, Guilin 541004, P. R. China*

⁹ *GuangXi University, Nanning 530004, P.R.China*

¹⁰ *Hangzhou Normal University, Hangzhou 310036, P. R. China*

¹¹ *Helmholtz Institute Mainz, J.J. Becherweg 45, D 55099 Mainz, Germany*

¹² *Henan Normal University, Xinxiang 453007, P. R. China*

¹³ *Henan University of Science and Technology, Luoyang 471003, P. R. China*

¹⁴ *Huangshan College, Huangshan 245000, P. R. China*

¹⁵ *Huazhong Normal University, Wuhan 430079, P. R. China*

¹⁶ *Hunan University, Changsha 410082, P. R. China*

¹⁷ *Indiana University, Bloomington, Indiana 47405, USA*

¹⁸ (A) *INFN Laboratori Nazionali di Frascati, Frascati, Italy;*

(B) *INFN and University of Perugia, I-06100, Perugia, Italy*

¹⁹ *Johannes Gutenberg University of Mainz,*

Johann-Joachim-Becher-Weg 45, 55099 Mainz, Germany

²⁰ *Joint Institute for Nuclear Research, 141980 Dubna, Russia*

²¹ *KVI/University of Groningen, 9747 AA Groningen, The Netherlands*

²² *Lanzhou University, Lanzhou 730000, P. R. China*

²³ *Liaoning University, Shenyang 110036, P. R. China*

²⁴ *Nanjing Normal University, Nanjing 210046, P. R. China*

²⁵ *Nanjing University, Nanjing 210093, P. R. China*

²⁶ *Nankai University, Tianjin 300071, P. R. China*

²⁷ *Peking University, Beijing 100871, P. R. China*

²⁸ *Seoul National University, Seoul, 151-747 Korea*

²⁹ *Shandong University, Jinan 250100, P. R. China*

³⁰ *Shanxi University, Taiyuan 030006, P. R. China*

³¹ *Sichuan University, Chengdu 610064, P. R. China*

³² *Soochow University, Suzhou 215006, P. R. China*

³³ *Sun Yat-Sen University, Guangzhou 510275, P. R. China*

- 88 ³⁴ *Tsinghua University, Beijing 100084, P. R. China*
89 ³⁵ (A)*Ankara University, Ankara, Turkey; (B)Dogus University,*
90 *Istanbul, Turkey; (C)Uludag University, Bursa, Turkey*
91 ³⁶ *Universitaet Giessen, 35392 Giessen, Germany*
92 ³⁷ *University of Hawaii, Honolulu, Hawaii 96822, USA*
93 ³⁸ *University of Minnesota, Minneapolis, MN 55455, USA*
94 ³⁹ *University of Rochester, Rochester, New York 14627, USA*
95 ⁴⁰ *University of Science and Technology of China, Hefei 230026, P. R. China*
96 ⁴¹ *University of South China, Hengyang 421001, P. R. China*
97 ⁴² *University of the Punjab, Lahore-54590, Pakistan*
98 ⁴³ (A)*University of Turin, Turin, Italy; (B)University of*
99 *Eastern Piedmont, Alessandria, Italy; (C)INFN, Turin, Italy*
100 ⁴⁴ *Wuhan University, Wuhan 430072, P. R. China*
101 ⁴⁵ *Zhejiang University, Hangzhou 310027, P. R. China*
102 ⁴⁶ *Zhengzhou University, Zhengzhou 450001, P. R. China*
- ^a *also at the Moscow Institute of Physics and Technology, Moscow, Russia*
^b *on leave from the Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine*
^c *also at the PNPI, Gatchina, Russia*
^d *now at Nagoya University, Nagoya, Japan*

Abstract

Hadronic transitions of $\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-$ ($J = 0, 1, 2$) are searched for using a sample of 1.06×10^8 $\psi(3686)$ events collected with the BESIII detector at the BEPCII storage ring. The η_c is reconstructed with $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ final states. No signals are observed in any of the three χ_{cJ} states in either η_c decay mode. At the 90% confidence level, the upper limits are determined to be $\mathcal{B}(\chi_{c0} \rightarrow \eta_c \pi^+ \pi^-) < 0.07\%$, $\mathcal{B}(\chi_{c1} \rightarrow \eta_c \pi^+ \pi^-) < 0.32\%$, and $\mathcal{B}(\chi_{c2} \rightarrow \eta_c \pi^+ \pi^-) < 0.54\%$. The upper limit of $\mathcal{B}(\chi_{c1} \rightarrow \eta_c \pi^+ \pi^-)$ is lower than the existing theoretical prediction by almost an order of magnitude. The branching fractions of $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$, $K^+ K^- \pi^+ \pi^- \pi^0$, $\omega K^+ K^-$ and $\phi \pi^+ \pi^- \pi^0$ ($J = 0, 1, 2$) are measured for the first time.

PACS numbers: 13.20.Gd, 13.25.Gv, 14.40.Pq

I. INTRODUCTION

Heavy quarkonia, both $c\bar{c}$ and $b\bar{b}$ bound states, have provided good laboratories for the study of the strong interaction [1, 2]. For the hadronic transitions between the heavy quarkonium states, Yan [3] characterized it as the emission of two soft gluons from the heavy quarks and the conversion of gluons into light hadrons. Based on this scheme, a series of decay rates, such as the E1-E1 hadronic transition, E1-M1 hadronic transition, M1-M1 hadronic transition have been calculated [4, 5]. It has been shown that the multipole expansion can make quite successful predictions for many hadronic transitions between the heavy quarkonia [1, 6]. However, most of these studies are for the transitions among the $^3S_1$ states; the hadronic transitions of ^3P_J states are seldom explored. Using a sample of $\Upsilon(3S)$, CLEO measured for the first time the transition rate of P -wave bottomonium $\chi_{bJ}(2P) \rightarrow \chi_{bJ}(1P)\pi\pi$ [7], and the results are consistent with the theoretical predictions [5]. For the hadronic transition of the P -wave charmonium states, there is only an upper limit of 2.2% at the 90% confidence level (C.L.) on the $\chi_{c2} \rightarrow \eta_c\pi^+\pi^-$ transition rate recently reported by the BaBar experiment [8]. The most promising process $\chi_{c1} \rightarrow \eta_c\pi\pi$, which is dominated by an E1-M1 transition, is calculated in the multipole expansion formalism, and a transition rate $\mathcal{B}(\chi_{c1} \rightarrow \eta_c\pi\pi) = (2.72 \pm 0.39)\%$ is predicted [9].

In this article, we search for $\chi_{cJ} \rightarrow \eta_c\pi^+\pi^-$ with η_c decays into $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$, where the K_S^0 is reconstructed in $\pi^+\pi^-$ and π^0 in $\gamma\gamma$ final states. We also report the first measurement of the branching fractions of $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+\pi^-$, $K^+ K^- \pi^+\pi^-\pi^0$, $\omega K^+ K^-$, and $\phi\pi^+\pi^-\pi^0$.

II. THE EXPERIMENT AND DATA SETS

The data sample for this analysis consists of 1.06×10^8 events produced at the peak of the $\psi(3686)$ resonance [10]; an additional 42 pb^{-1} of data were collected at a center-of-mass energy of $\sqrt{s}=3.65 \text{ GeV}$ to determine non-resonant continuum background contributions. The data are accumulated with the BESIII detector operated at the BEPCII e^+e^- collider.

The BESIII detector, described in detail in Ref. [11], has an effective geometrical acceptance of 93% of 4π . It contains a small cell helium-based main drift chamber (MDC) which provides momentum measurements of charged particles; a time-of-flight system (TOF) based on plastic scintillator which helps to identify charged particles; an electromagnetic calorimeter (EMC) made of CsI (Tl) crystals which is used to measure the energies of photons and provide trigger signals; and a muon system (MUC) made of Resistive Plate Chambers (RPC). The resolution of the charged particles is 0.5% at 1 GeV/ c in a 1 Tesla magnetic field. The energy loss (dE/dx) measurement provided by the MDC has a resolution better than 6% for electrons from Bhabha scattering. The photon energy resolution can reach 2.5% (5%) at 1 GeV in the barrel (endcaps) of the EMC. And the time resolution of TOF is 80 ps in the barrel and 110 ps in the endcaps.

Monte Carlo (MC) simulated events are used to determine the detection efficiency, optimize the selection criteria, and study the possible backgrounds. The simulation of the BESIII detector is GEANT4 [12] based, where the interactions of the particles with the detector material are simulated. The $\psi(3686)$ resonance is produced with KKMC [13], while the subsequent decays are generated with EVTGEN [14]. The study of the background is based on a sample of 10^8 $\psi(3686)$ inclusive decays which are generated with known branching fractions taken from the Particle Data Group (PDG) [15], or with LUNDCHARM [16] for

the unmeasured decays.

III. EVENT SELECTION

A charged track should have good quality in the track fitting and be within the angle coverage of the MDC, $|\cos\theta| < 0.93$. A good charged track (excludes those from K_S^0 decays) is required to be within 1 cm of the e^+e^- annihilation interaction point (IP) transverse to the beam line and within 10 cm of the IP along the beam axis. Charged-particle identification (PID) is based on combining the dE/dx and TOF information in the variable $\chi_{\text{PID}}^2(i) = (\frac{dE/dx_{\text{measured}} - dE/dx_{\text{expected}}}{\sigma_{dE/dx}})^2 + (\frac{\text{TOF}_{\text{measured}} - \text{TOF}_{\text{expected}}}{\sigma_{\text{TOF}}})^2$. The values $\chi_{\text{PID}}^2(i)$ and the corresponding confidence levels $\text{Prob}_{\text{PID}}(i)$ are calculated for each charged track for each particle hypothesis i (pion, kaon, or proton).

Photons are reconstructed from isolated showers in the EMC which are at least 20 degrees away from any of the charged tracks. In order to improve the reconstruction efficiency and the energy resolution, the energy deposited in the nearby TOF counter is included. Photon candidates are required to have the energy greater than 25 MeV in the EMC barrel region ($|\cos\theta| < 0.8$), while in the EMC endcap region ($0.86 < |\cos\theta| < 0.92$), the energy threshold requirement is increased to 50 MeV. EMC timing requirements are used to suppress noise and energy deposits unrelated to the event.

K_S^0 candidates are reconstructed from secondary vertex fits to all the charged-track pairs in an event (assume the tracks to be π). The combination with the best fit quality is kept, and the K_S^0 candidate must have an invariant mass within 7 MeV/ c^2 of the K_S^0 nominal mass and the secondary vertex be at least 0.5 cm away from the IP. The reconstructed K_S^0 information is used as input for the subsequent kinematic fit. The π^0 candidates are reconstructed from pairs of photons with an invariant mass in the range $0.120 \text{ GeV}/c^2 < M(\gamma\gamma) < 0.145 \text{ GeV}/c^2$.

In selecting $\psi(3686) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$, a candidate event should have at least six charged tracks and at least one good photon. After K_S^0 selection, the event should have exactly four additional good charged tracks with zero net charge. While in selecting $\psi(3686) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$, a candidate event should have four good charged tracks with zero net charge and at least three good photons. The $\gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($\gamma K^+ K^- \pi^+ \pi^- \pi^0$) candidate is then subjected to a four-constraint (4C) kinematic fit to reduce background and improve the mass resolution. Determination of the species of the final state particles and selection of the best photons when additional photons (and π^0 candidates) are found in an event are achieved by selecting the combination with the minimum value of $\chi^2 = \chi_{4C}^2 + \sum_{j=1}^4 \chi_{\text{PID}}^2(j)$, where χ_{4C}^2 is the chi-square from the 4C kinematic fit. Events with $\chi_{4C}^2 < 50$ are kept as $\gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($\gamma K^+ K^- \pi^+ \pi^- \pi^0$) candidates.

There is substantial background from $\psi(3686) \rightarrow X + J/\psi$ decays. The $\psi(3686) \rightarrow \pi^+ \pi^- J/\psi$ events are removed by requiring the recoil mass of any $\pi^+ \pi^-$ pair to be outside of a $\pm 3\sigma$ window around the J/ψ nominal mass, where σ is the resolution of the $\pi^+ \pi^-$ recoil mass. The $\psi(3686) \rightarrow \eta J/\psi$, $\eta \rightarrow \gamma \pi^+ \pi^-$ events are rejected if $0.535 \text{ GeV}/c^2 < M(\gamma \pi^+ \pi^-) < 0.555 \text{ GeV}/c^2$. To reject background with one more photon than the signal events ($\gamma \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ or $\gamma \gamma K^+ K^- \pi^+ \pi^- \pi^0$), $\chi_{4C}^2 > 10$ is required when a 4C kinematic fit to $\gamma \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ or $\gamma \gamma K^+ K^- \pi^+ \pi^- \pi^0$ is applied to the event; and to suppress $\psi(3686) \rightarrow \pi^0 \pi^0 J/\psi$, $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$ background, $M(K^+ K^- \pi^+ \pi^-) > 3.125 \text{ GeV}/c^2$ or $M(K^+ K^- \pi^+ \pi^-) < 3.095 \text{ GeV}/c^2$ is required.

IV. DATA ANALYSIS

A. $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ and $\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

After the above selection, the invariant mass distributions of the hadron system are shown in Fig. 1, and clear χ_{cJ} signals are observed with very low background level in the two modes.




FIG. 1: Invariant mass spectrum of $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ (left panel) and $K^+ K^- \pi^+ \pi^- \pi^0$ (right panel), together with the best fit results. The points with error bars are data, and the solid lines are the total fit results. The χ_{c0} , χ_{c1} , and χ_{c2} signals are shown as dotted lines, dash-dotted lines, and long-dashed lines, respectively; the backgrounds are in dashed lines.

Using the inclusive MC events sample, the potential backgrounds from the $\psi(3686)$ decays which may contaminate $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$) are estimated. Events from $\psi(3686) \rightarrow \gamma \chi_{cJ}$, $\chi_{cJ} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$ and $\chi_{c0} \rightarrow K_S^0 K_S^0 K^+ K^-$ ($\psi(3686) \rightarrow \gamma \chi_{cJ}$, $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^0$) may peak at the signal region, and the contributions from these peaking backgrounds are estimated using the detection efficiencies determined from the MC simulations and the corresponding branching fractions from previous measurements [15], and then subtracted. These background events after final event selection are listed in Table I and II; the errors in the numbers of events are from the uncertainties of the detection efficiency and branching fractions. The other backgrounds are composed of dozens of decay modes and smoothly distributed in the full mass region ($3.3 \text{ GeV}/c^2 \sim 3.6 \text{ GeV}/c^2$); this kind of background is described by a second-order Chebyshev function.

TABLE I: The number of remanent peaking background events ($N_{\text{bkg}}^{\text{peak}}$) in $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ after final event selection. The branching fractions ($\mathcal{B}$) are taken from PDG [15].

Decay modes	$N_{\text{bkg}}^{\text{peak}}$	$\mathcal{B}$ [15]
$\chi_{c0} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	4.9 ± 1.4	$(5.8 \pm 1.1) \times 10^{-3}$
$\chi_{c1} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	0.1 ± 0.1	$(7.2 \pm 3.1) \times 10^{-4}$
$\chi_{c2} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	0.6 ± 0.3	$(2.4 \pm 0.6) \times 10^{-3}$
$\chi_{c0} \rightarrow K_S^0 K_S^0 K^+ K^-$	43.3 ± 15.8	$(1.4 \pm 0.5) \times 10^{-3}$

TABLE II: The number of remanent peaking background events ($N_{\text{bkg}}^{\text{peak}}$) in $\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ after final event selection. The branching fractions ($\mathcal{B}$) are taken from PDG [15].

Decay modes	$N_{\text{bkg}}^{\text{peak}}$	$\mathcal{B}$ [15]
$\chi_{c0} \rightarrow K_S^0 K^\pm \pi^\mp \pi^0$	123.8 ± 19.4	$\frac{1}{2} \times (2.52 \pm 0.34) \times 10^{-2}$
$\chi_{c1} \rightarrow K_S^0 K^\pm \pi^\mp \pi^0$	39.0 ± 9.1	$\frac{1}{2} \times (9.0 \pm 1.5) \times 10^{-3}$
$\chi_{c2} \rightarrow K_S^0 K^\pm \pi^\mp \pi^0$	52.5 ± 9.3	$\frac{1}{2} \times (1.51 \pm 0.22) \times 10^{-2}$

Data taken at $\sqrt{s} = 3.65$ GeV are used to estimate backgrounds from the continuum process $e^+e^- \rightarrow q\bar{q}$. This kind of background is found to be small and uniformly distributed in the full mass region of interest in both decay modes, so the contribution can be represented by the smooth background term in the fit.

In the measurement of $\mathcal{B}(\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)$, the contributions from intermediate states with narrow resonances such as η , ω , and ϕ are excluded; the identification of such resonances is similar to that used in Refs. [19, 20]. The branching fractions of these decays $\chi_{cJ} \rightarrow \eta K^+ K^-$, $\chi_{cJ} \rightarrow \omega K^+ K^-$, $\chi_{cJ} \rightarrow \phi K^+ K^-$, and $\chi_{cJ} \rightarrow \phi \pi^+ \pi^- \pi^0$ are measured using the same data sample. The fits to the invariant mass spectrum of $\pi^+ \pi^- \pi^0$ and $K^+ K^-$ in the three χ_{cJ} signal regions (as defined in Sect. IV B)) are shown in Fig. 2, and the results are listed in Table III. The first errors are statistical and the second ones are systematic. The sources of the systematic errors are similar to those in the measurement of $\mathcal{B}(\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)$, as will be shown in Sect. V. The branching fractions of $\chi_{cJ} \rightarrow \omega K^+ K^-$ and $\chi_{cJ} \rightarrow \phi \pi^+ \pi^- \pi^0$ are the first measurement, and those of other modes from this measurement are consistent within errors with the known PDG values [15] when available. Events from these decay modes are removed by requiring $|m_{\pi^+ \pi^- \pi^0} - m_\eta| > 15 \text{ MeV}/c^2$, $|m_{\pi^+ \pi^- \pi^0} - m_\omega| > 40 \text{ MeV}/c^2$, $|m_{\pi^+ \pi^- \pi^0} - m_\phi| > 15 \text{ MeV}/c^2$ and $|m_{K^+ K^-} - m_\phi| > 15 \text{ MeV}/c^2$. The contribution from $\chi_{cJ} \rightarrow \phi\phi$, $\phi \rightarrow K^+ K^-$, $\phi \rightarrow \pi^+ \pi^- \pi^0$ events is also removed by these requirements. The expected remaining events from these decay channels are also listed in Table III and will be subtracted from the signal yields from the best fits.

An unbinned maximum likelihood fit is applied to the invariant mass spectrum of $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($K^+ K^- \pi^+ \pi^- \pi^0$) to extract the numbers of χ_{cJ} events in Fig. 1. The χ_{cJ} signals are described by the corresponding MC simulated signal shape convolved with a Gaussian function $G(\mu, \sigma)$ to take into account the difference in the mass scale and the mass resolution between data and MC simulation. The means (μ) and the standard deviations (σ) of the Gaussian functions are floated parameters in the fit. In the generation of the χ_{c0} MC events, the E1 radiative transition factor E_γ^3 is included, where E_γ is the energy of the radiative photon in the $\psi(3686)$ rest frame. To damp the diverging tail due to the E_γ^3 dependence, a damping function $\frac{E_0^2}{E_\gamma E_0 + (E_\gamma - E_0)^2}$ used by KEDR [18] is introduced, where $E_0 = (m_{\psi(3686)}^2 - m_{\chi_{c0}}^2)/2m_{\psi(3686)}$. The backgrounds are described by a second-order Chebyshev function in both decay modes.

The fit to the invariant mass spectrum of $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ yields 2837 ± 64 , 5180 ± 75 , and 4560 ± 71 signal events for χ_{c0} , χ_{c1} , and χ_{c2} , respectively; while the fit to the $K^+ K^- \pi^+ \pi^- \pi^0$ modes yields 9372 ± 130 , 12415 ± 126 , and 11366 ± 123 events for χ_{c0} , χ_{c1} , and χ_{c2} , respectively.




FIG. 2: Fits to the invariant mass spectrum of $\pi^+\pi^-\pi^0$ (left panels) and K^+K^- (right panels) in three χ_{cJ} mass region after $\chi_{cJ} \rightarrow \phi\phi$, $\phi \rightarrow K^+K^-$, $\phi \rightarrow \pi^+\pi^-\pi^0$ were rejected. From top to bottom are in χ_{c0} mass region, χ_{c1} mass region, and χ_{c2} mass region, respectively.

B. $\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-$

To study the $\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-$ transitions, we define the signal regions for χ_{c0} , χ_{c1} , and χ_{c2} as $3.38 \text{ GeV}/c^2 < M(K\bar{K}\pi\pi\pi) < 3.45 \text{ GeV}/c^2$, $3.48 \text{ GeV}/c^2 < M(K\bar{K}\pi\pi\pi) < 3.54 \text{ GeV}/c^2$, and $3.54 \text{ GeV}/c^2 < M(K\bar{K}\pi\pi\pi) < 3.60 \text{ GeV}/c^2$, respectively. Here, $K\bar{K}\pi\pi\pi$ is either $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ or $K^+ K^- \pi^+ \pi^- \pi^0$. The invariant mass spectra of $K_S^0 K^\pm \pi^\mp$ ($K^+ K^- \pi^0$) with $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($K^+ K^- \pi^+ \pi^- \pi^0$) in the three χ_{cJ} signal regions are shown in Fig. 3. In both decay modes, there are no significant η_c signal in χ_{c0} and χ_{c1} decay; the η_c signal observed in χ_{c2} decays is found to be mainly from $\psi(3686) \rightarrow \pi^+ \pi^- J/\psi$, $J/\psi \rightarrow \gamma \eta_c$, $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ ($\eta_c \rightarrow K^+ K^- \pi^0$), as described below.

The potential backgrounds from $\psi(3686)$ decays are investigated with the inclusive

TABLE III: The number of background events ($N_{\text{bkg}}^{\text{peak}}$) remained in $\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ from modes with narrow intermediate states. The branching fraction of $\chi_{cJ} \rightarrow \phi\phi$ is taken from the BESIII measurement [19] with both statistical and systematic errors; the other branching fractions are measured in this analysis. We also list PDG values [15] in the last column for comparison.

Decay modes	$N_{\text{bkg}}^{\text{peak}}$	Branching fraction	PDG [15] value
$\chi_{c0} \rightarrow \eta K^+ K^-$	0	forbidden by J^P -conservation	$< 2.3 \times 10^{-4}$
$\chi_{c1} \rightarrow \eta K^+ K^-$	3.81 ± 0.70	$(3.48 \pm 0.57) \times 10^{-4}$	$(3.3 \pm 1.0) \times 10^{-4}$
$\chi_{c2} \rightarrow \eta K^+ K^-$	1.90 ± 0.53	$(1.69 \pm 0.45) \times 10^{-4}$	$< 3.5 \times 10^{-4}$
$\chi_{c0} \rightarrow \omega K^+ K^-$	136.87 ± 10.84	$(1.94 \pm 0.06 \pm 0.20) \times 10^{-3}$	-
$\chi_{c1} \rightarrow \omega K^+ K^-$	69.24 ± 6.20	$(7.82 \pm 0.36 \pm 0.84) \times 10^{-4}$	-
$\chi_{c2} \rightarrow \omega K^+ K^-$	57.26 ± 5.04	$(7.32 \pm 0.39 \pm 0.78) \times 10^{-4}$	-
$\chi_{c0} \rightarrow \phi K^+ K^-$	30.37 ± 4.91	$(1.15 \pm 0.17) \times 10^{-3}$	$(0.98 \pm 0.25) \times 10^{-3}$
$\chi_{c1} \rightarrow \phi K^+ K^-$	17.61 ± 3.79	$(6.74 \pm 1.37) \times 10^{-4}$	$(4.3 \pm 1.6) \times 10^{-4}$
$\chi_{c2} \rightarrow \phi K^+ K^-$	30.39 ± 5.27	$(1.14 \pm 0.16) \times 10^{-3}$	$(1.55 \pm 0.33) \times 10^{-3}$
$\chi_{c0} \rightarrow \phi \pi^+ \pi^- \pi^0$	49.57 ± 2.68	$(1.18 \pm 0.07 \pm 0.13) \times 10^{-3}$	-
$\chi_{c1} \rightarrow \phi \pi^+ \pi^- \pi^0$	28.59 ± 1.99	$(7.54 \pm 0.53 \pm 0.80) \times 10^{-4}$	-
$\chi_{c2} \rightarrow \phi \pi^+ \pi^- \pi^0$	34.45 ± 2.36	$(9.25 \pm 0.63 \pm 0.97) \times 10^{-4}$	-
$\chi_{c0} \rightarrow \phi\phi$	0.97 ± 0.17	$(8.0 \pm 0.3 \pm 0.8) \times 10^{-4}$	$(8.2 \pm 0.8) \times 10^{-4}$
$\chi_{c1} \rightarrow \phi\phi$	0.38 ± 0.08	$(4.4 \pm 0.3 \pm 0.5) \times 10^{-4}$	-
$\chi_{c2} \rightarrow \phi\phi$	1.03 ± 0.20	$(10.7 \pm 0.3 \pm 1.2) \times 10^{-4}$	$(11.4 \pm 1.2) \times 10^{-4}$

MC events. The dominant backgrounds are the irreducible $\psi(3686) \rightarrow \gamma\chi_{cJ}$, $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$) events. These events have the same final states as the signal events but $K_S^0 K^\pm \pi^\mp$ ($K^+ K^- \pi^0$) are not from the decay of η_c . In the χ_{c2} signal region, there are peaking backgrounds from the decay $\psi(3686) \rightarrow \pi^+ \pi^- J/\psi$, $J/\psi \rightarrow \gamma\eta_c$, $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ ($\eta_c \rightarrow K^+ K^- \pi^0$). The energy of the transition photon in this decay is close to the energy of the transition photon in $\psi(3686) \rightarrow \gamma\chi_{c2}$, and the final states are the same as the signal events. The other backgrounds are composed of dozens of decay channels, each with a small contribution. The dominant backgrounds and the other backgrounds contribute a smooth component in the η_c mass region ($2.7 \text{ GeV}/c^2 \sim 3.2 \text{ GeV}/c^2$), so these backgrounds are described by a second-order Chebyshev function as shown in Fig. 3. In the χ_{c2} case, the peaking background has the same final state and similar kinematics as the signal events, so we use the same line-shape to describe both of them. Data taken at $\sqrt{s} = 3.65 \text{ GeV}$ are used to estimate backgrounds from the continuum process $e^+ e^- \rightarrow q\bar{q}$. It is found that this background is small and uniformly distributed in the full mass region in both decay modes, so the contribution is neglected.

An unbinned maximum likelihood fit is applied to the invariant mass spectrum of $K_S^0 K^\pm \pi^\mp$ ($K^+ K^- \pi^0$) to extract the number of η_c events, as shown in Fig. 3. The η_c signal is described by a MC simulated line shape with the detector resolution included, and the resonance parameters of η_c are fixed to the latest measurement from the BESIII experiment [17]. The background (except the peaking background in the χ_{c2} signal region) is described with a second-order Chebyshev polynomial function in both decay modes in the three χ_{cJ} signal regions.




FIG. 3: Invariant mass spectra of $K_S^0 K^\pm \pi^\mp$ (top row) and $K^+ K^- \pi^0$ (bottom row) with $K \bar{K} \pi \pi$ in χ_{c0} (left panel), χ_{c1} (middle panel), and χ_{c2} (right panel) signal regions and the fit results. Dots with error bars are data; the solid lines are the total results from the best fits to the invariant mass spectrum. The η_c signals are shown in dash-dotted lines (in χ_{c2} mass region, the contribution from the peaking background is not removed.); the backgrounds as dashed lines.

As there is no significant η_c signal in any of the three χ_{cJ} states in either η_c decay mode, we set upper limits on $\mathcal{B}(\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-)$ using the probability density function (PDF) for the expected number of signal events. In the χ_{c0} and χ_{c1} signal regions, the likelihood distributions in the fitting of the invariant mass spectra in Fig. 3 are taken as the PDFs directly. They are obtained by setting the number of η_c signal events from zero up to a very large number. In the χ_{c2} signal region, the likelihood distribution also contains the contribution from the peaking background. Using the known branching fractions [15], the detection efficiency from MC simulation, and the number of $\psi(3686)$ events, the expected peaking background are 45.7 ± 11.6 in $K_S^0 K^\pm \pi^\mp$ and 34.4 ± 8.7 in $K^+ K^- \pi^0$. Here, the errors include the uncertainties in the detection efficiency and the branching fractions. Then the PDF of signal is extracted with the PDF of the peaking background (Gaussian distribution with mean set to the expected number of peaking backgrounds, sigma set to its error) and the PDF from the fit. The systematic uncertainties are considered by smearing the PDF in each decay with a Gaussian. The upper limit on the number of events at the 90% C.L. is defined as N^{up} , corresponding to the number of events at 90% of the integral of the smeared PDF. In each decay mode in the three χ_{cJ} states, the fit-related systematic errors on the number of signal yield are estimated by using different fit ranges, different orders of the background polynomial, and different η_c line shapes with the parameters of η_c changed by one standard deviation [17]; the maximum N^{up} is used in the upper limit calculation.

V. SYSTEMATIC UNCERTAINTIES

The systematic uncertainties in the measurement of $\mathcal{B}(\chi_{cJ} \rightarrow K\bar{K}\pi\pi\pi)$ and $\mathcal{B}(\chi_{cJ} \rightarrow \eta_c\pi^+\pi^-)$ are summarized in Tables IV and V, respectively. The systematic errors related to the MDC tracking (2% per track for those from IP), photon reconstruction (1% per photon), and π^0 reconstruction (1%) are estimated with control samples [20, 21]; the errors in the branching fractions of $\psi(3686) \rightarrow \gamma\chi_{cJ}$ and $\eta_c \rightarrow K\bar{K}\pi$ are taken from the PDG [15] and are propagated to the χ_{cJ} branching fraction measurement; and a 2% uncertainty is taken for each decay due to the limited statistics of the MC samples used. There is an overall 4% uncertainty in the branching fraction associated with the determination of the number of $\psi(3686)$ events in our data sample [10].

TABLE IV: Systematic errors (in %) in $\mathcal{B}(\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)$ and $\mathcal{B}(\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)$.

Sources	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$			$K^+ K^- \pi^+ \pi^- \pi^0$		
	χ_{c0}	χ_{c1}	χ_{c2}	χ_{c0}	χ_{c1}	χ_{c2}
MDC tracking	8.0			8.0		
Photon reconstruction	1.0			3.0		
MC statistics	1.1	1.4	1.5	1.0	1.4	1.4
K_S^0 reconstruction	1.4	1.6	1.7	—	—	—
π^0 reconstruction	—	—	—	1.0		
Kinematic fit	1.5	1.9	1.7	0.4	0.4	0.2
Damping function	0.5	0.1	0.1	0.4	0.1	0.1
Intermediate states	1.0	1.0	1.0	4.0	4.0	4.0
Fitting range	1.0	0.4	0.2	0.4	0.4	0.7
Background shape	1.4	0.7	0.6	1.3	0.7	0.4
$\mathcal{B}(\psi(3686) \rightarrow \gamma\chi_{cJ})$	3.2	4.4	3.9	3.2	4.4	3.9
Number of $\psi(3686)$ events	4.0			4.0		
Total	10.1	10.5	10.3	10.9	11.3	11.2

A. K_S^0 reconstruction

The uncertainty in the K_S^0 reconstruction arises from three parts: the geometric acceptance, the tracking efficiency, and the efficiency of K_S^0 selection. The first part is estimated using MC simulation, and the other two are studied using $J/\psi \rightarrow K^{*\pm} K^\mp$, $K^{*\pm} \rightarrow K^0 \pi^\pm$. By selecting a pair of $K^\pm \pi^\mp$, the recoil mass spectrum shows a clear K^0 signal. The efficiency of K_S^0 reconstruction is calculated with $\frac{n_1}{\mathcal{B}(K_S^0 \rightarrow \pi^+ \pi^-) \times (n_1 + n_2)/2}$, where n_1 is the number of K^0 obtained from a fit to the $K^\pm \pi^\mp$ recoiling mass when there is a K_S^0 reconstructed in the recoil side that satisfies the K_S^0 selection, and n_2 is the number of K^0 from fitting to the $K^\pm \pi^\mp$ recoiling mass spectrum when no K_S^0 candidate satisfies the K_S^0 selection. The difference in the efficiency of K_S^0 reconstruction ($\varepsilon^{\text{data}}/\varepsilon^{\text{MC}} - 1$) as a function of K_S^0 momentum is shown in Fig. 4. The difference in the K_S^0 reconstruction between data and MC simulation is fitted with a linear function of the K_S^0 momentum as shown in Fig. 4 together with the

TABLE V: Systematic errors (in %) in $\mathcal{B}(\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-)$ in $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ and $\eta_c \rightarrow K^+ K^- \pi^0$ decay modes.

Sources	$\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$			$\eta_c \rightarrow K^+ K^- \pi^0$		
	χ_{c0}	χ_{c1}	χ_{c2}	χ_{c0}	χ_{c1}	χ_{c2}
MDC tracking	8.0			8.0		
Photon reconstruction	1.0			3.0		
MC statistics	1.8	1.5	1.6	1.8	1.5	1.7
K_S^0 reconstruction	2.4	2.3	2.2	—	—	—
π^0 reconstruction	—	—	—	1.0		
Kinematic fit	1.5	1.7	1.8	0.9	0.2	0.4
$\mathcal{B}(\psi(3686) \rightarrow \gamma \chi_{cJ})$	3.2	4.4	3.9	3.2	4.4	3.9
$\mathcal{B}(\eta_c \rightarrow K \bar{K} \pi)$	8.4			8.4		
Number of $\psi(3686)$ events	4.0			4.0		
Total	13.2	13.5	13.4	13.2	13.5	13.4

$\pm 1\sigma$ envelopes. Since the difference between data and MC is significant, we do a correction to the signal MC according to the momentum of K_S^0 , and the uncertainty of this correction is taken as the systematic error.

The systematic errors in $\mathcal{B}(\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)$ are found to be 1.4%, 1.6%, and 1.7%, for $J = 0, 1$, and 2, respectively; while for $\mathcal{B}(\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-)$, they are 2.4%, 2.3%, and 2.2% for χ_{c0} , χ_{c1} , and χ_{c2} , respectively.




FIG. 4: The difference in the K_S^0 reconstruction efficiency between data and MC simulation (points with error bars), together with the fit to the difference with a linear function of momentum. The solid line is from the best fit and the dashed lines are the $\pm 1\sigma$ envelopes of the best fit.

B. Kinematic fit

In the MC simulation, the model is much simpler than the real detector performance, and this results in differences between data and MC simulation in the track parameters of photons and charged tracks. The simulation of the photon has been checked in another analysis [10],

which shows good agreement between data and MC simulation. For the charged tracks, careful comparisons with purely selected data samples indicate that the MC simulates the momentum and angular resolutions significantly better than those in data, while the error matrix elements agree well between data and MC simulation. This results in a much narrower χ_{4C}^2 distribution in MC than in data, and introduces a bias in the efficiency estimation. We correct the track helix parameters of MC simulation to reduce the difference between data and MC simulation.

We use $J/\psi \rightarrow \phi f_0(980)$, $\phi \rightarrow K^+ K^-$, $f_0(980) \rightarrow \pi^+ \pi^-$ as a control sample to study the difference on the helix parameters of charged tracks between data and MC, as this channel has a large production rate, very low background, and has both pions and kaons. We find that the pull distributions of data are wider than MC simulation and the peak positions are shifted. These obvious differences between data and MC suggest wrong track parameters have been set in MC simulation. The helix parameters of each track in the MC simulation are enlarged by smearing with a Gaussian function $G(\mu, \sigma) = G((\mu_i^{\text{data}} - \mu_i^{\text{MC}}) \times V_{ii}, \sqrt{(\sigma_i^{\text{data}}/\sigma_i^{\text{MC}})^2 - 1} \times V_{ii})$, where $i = \{d\rho, \phi_0, \kappa, dz, tg\lambda\}$ is the i -th helix parameter of the track and V is the corresponding covariance matrix. Here $d\rho$ is the distance from the pivot to the orbit in the x - y plane, ϕ_0 is the azimuthal angle specifies the pivot with respect to the helix center, κ is the reciprocal of the transverse momentum, dz is the distance of the helix from the pivot to the orbit in the z direction, and $tg\lambda$ is the slope of the track. The correction factors μ_i^{data} , μ_i^{MC} , σ_i^{data} , and σ_i^{MC} are the means and resolutions of the pull distributions of the data and MC obtained from control samples.

The correction factors are listed in Table VI. If the correction is perfect, χ_{4C}^2 distributions of data and MC simulation will be consistent with each other; however, from the comparisons of many final states, we find that the agreement between data and MC simulation does improve significantly but differences still exist. This indicates that the effect is from multiple sources, and our procedure cannot solve all the problems. In our analysis, we take the efficiency from the track-parameter-corrected MC samples as the nominal value, and take half of the difference between MC samples before and after the correction as the systematic error from the kinematic fitting. This is a very conservative estimation. The comparison of χ_{4C}^2 distributions between data and MC simulation before and after the track-parameter-correction are shown in Figs. 5 and 6 for $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ and $K^+ K^- \pi^+ \pi^- \pi^0$, respectively.

TABLE VI: Correction factors extracted from pull distributions using a control sample of $J/\psi \rightarrow \phi f_0(980)$, $\phi \rightarrow K^+ K^-$, $f_0(980) \rightarrow \pi^+ \pi^-$.

	ϕ_0		κ		$tg\lambda$	
	$\mu^{\text{data}} - \mu^{\text{MC}}$	$\sigma^{\text{data}}/\sigma^{\text{MC}}$	$\mu^{\text{data}} - \mu^{\text{MC}}$	$\sigma^{\text{data}}/\sigma^{\text{MC}}$	$\mu^{\text{data}} - \mu^{\text{MC}}$	$\sigma^{\text{data}}/\sigma^{\text{MC}}$
K^+	-0.04 ± 0.03	1.19 ± 0.02	-0.24 ± 0.03	1.28 ± 0.02	-0.38 ± 0.01	1.25 ± 0.02
K^-	0.06 ± 0.03	1.21 ± 0.02	0.25 ± 0.03	1.25 ± 0.02	-0.36 ± 0.01	1.21 ± 0.02
π^+	-0.06 ± 0.03	1.25 ± 0.02	-0.10 ± 0.03	1.31 ± 0.02	-0.36 ± 0.01	1.25 ± 0.02
π^-	-0.02 ± 0.03	1.23 ± 0.02	0.10 ± 0.03	1.27 ± 0.02	-0.36 ± 0.01	1.21 ± 0.02

The systematic errors in $\mathcal{B}(\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)$ ($\mathcal{B}(\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)$) are 1.5%, 1.9%, and 1.7% (0.4%, 0.4%, and 0.2%) for $J = 0, 1$, and 2 , respectively; and 1.5%, 1.7%,




FIG. 5: Comparison of χ^2_{4C} between signal MC and data for $\psi(3686) \rightarrow \gamma\chi_{cJ}$, $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$. The points with error bars are data, and the solid lines are MC simulation. Left panel: signal MC without track-parameter-correction; Right panel: signal MC after track-parameter-correction. From top to bottom are χ_{c0} , χ_{c1} , and χ_{c2} , respectively.

and 1.8% (0.9%, 0.2%, and 0.4%) systematic uncertainties are assigned to $\mathcal{B}(\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-)$ with $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ ($K^+ K^- \pi^0$) for $J = 0, 1$, and 2, respectively.

C. Uncertainty from damping factor

In the fit to the invariant mass spectrum of $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ and $K^+ K^- \pi^+ \pi^- \pi^0$, the damping function used by KEDR is adopted. Another damping factor used by CLEO [22] is $e^{-E_\gamma^2/8\beta^2}$ with $\beta = (0.0650 \pm 0.0025)$ GeV determined from their fit. Using this damping function with $\beta = (0.097 \pm 0.024)$ GeV which is extracted from fitting χ_{c0} data, the differences




FIG. 6: Comparison of χ^2_{4C} between signal MC and data for $\psi(3686) \rightarrow \gamma\chi_{cJ}$, $\chi_{cJ} \rightarrow K^+K^-\pi^+\pi^-\pi^0$. The points with error bars are data, and the solid lines are signal MC plus possible background estimated using inclusive MC sample. Left panel: signal MC without track-parameter-correction; Right panel: signal MC after track-parameter-correction. From top to bottom are χ_{c0} , χ_{c1} , and χ_{c2} , respectively.

on the branching fractions of $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$) are assigned to the systematic error due to damping function, which are 0.5%, 0.1%, and 0.1% (0.4%, 0.1%, and 0.1%) for $J = 0, 1$, and 2 , respectively. The effect for χ_{c1} and χ_{c2} is small since the two states are very narrow.

D. Uncertainty from intermediate states

The detection efficiencies for the measurement of $\mathcal{B}(\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)$ and $\mathcal{B}(\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)$ are estimated using the MC simulation with χ_{cJ} decay to $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ and $K^+ K^- \pi^+ \pi^- \pi^0$ generated according to pure phase space distribution. From the data, we see broad intermediate states such as K^* and ρ in the invariant mass spectra of $K\pi$ and $\pi\pi$. The branching fractions of $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$) via these intermediate states are measured by fitting the invariant mass spectra of $K\pi$ and $\pi\pi$. An alternative signal MC sample is generated with all possible intermediate states and corresponding branching fractions to determine the efficiency. The efficiency difference between this sample and the phase space sample is about 1.0% for $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ and 4.0% for $\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$; these are taken as the systematic error due to intermediate states.

E. Uncertainty from fitting

The systematic uncertainty due to the fitting range is estimated by fitting the invariant mass spectrum in the range $3.25 \text{ GeV}/c^2 \sim 3.61 \text{ GeV}/c^2$ and $3.35 \text{ GeV}/c^2 \sim 3.60 \text{ GeV}/c^2$. The biggest differences in the branching fractions are assigned as errors, which are 1.0%, 0.4%, and 0.2% for χ_{c0} , χ_{c1} , and χ_{c2} , respectively, in $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ decay; and 0.4%, 0.4%, and 0.7% for χ_{c0} , χ_{c1} , and χ_{c2} , respectively, in $K^+ K^- \pi^+ \pi^- \pi^0$ decay. The background shape is changed from a second-order Chebyshev polynomial function to a third-order Chebyshev polynomial function, and the differences are taken to be the systematic errors, which are 1.4%, 0.7%, and 0.6% for χ_{c0} , χ_{c1} , and χ_{c2} , respectively, in $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$; and 1.3%, 0.7%, and 0.4% for χ_{c0} , χ_{c1} , and χ_{c2} in $K^+ K^- \pi^+ \pi^- \pi^0$.

VI. RESULTS AND DISCUSSION

Using the numbers of signal χ_{cJ} events from the fits, together with the corresponding efficiencies, the branching fractions of $\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ ($\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$) are determined and listed in Table VII. In the branching fractions of $\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$, contributions from narrow resonances η , ω , and ϕ are subtracted. All these are first measurements, and the branching fractions are at the 1% level. Comparing the two decay modes, we found the ratio of the branching fractions is around one-half which may be a consequence of isospin symmetry. We also measured the branching fractions of $\chi_{cJ} \rightarrow \omega K^+ K^-$ and $\chi_{cJ} \rightarrow \phi \pi^+ \pi^- \pi^0$ for the first time, the results are listed in Table VII also.

With the upper limit on the numbers of events at the 90% C.L. in $\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-$, $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ ($\eta_c \rightarrow K^+ K^- \pi^0$), as well as the corresponding efficiencies, the upper limits on the branching fraction of $\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-$ in the two decay modes are determined, as listed in Table VIII. We give a more stringent constraint on the $\chi_{c2} \rightarrow \eta_c \pi^+ \pi^-$ branching fraction than BaBar does [8]. The theoretical prediction of $\mathcal{B}(\chi_{c1} \rightarrow \eta_c \pi^+ \pi^-)$ is also listed in Table VIII, which is larger than our measurements. We note that the theoretical prediction uses experimental results as input to normalize the parameters in the model. For example, the parameter α_M/α_E is extracted by comparing the branching fraction of $\psi(3686) \rightarrow h_c \pi^0$ between the theoretical calculation and the experimental measurement. This makes the prediction highly dependent on the former experimental results and theoretical models.

TABLE VII: The results for $\mathcal{B}(\chi_{cJ} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)$, $\mathcal{B}(\chi_{cJ} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)$, $\mathcal{B}(\chi_{cJ} \rightarrow \omega K^+ K^-)$, and $\mathcal{B}(\chi_{cJ} \rightarrow \phi \pi^+ \pi^- \pi^0)$. The first errors are statistical and the second ones are systematic.

Decay mode	N^{signal}	ϵ (%)	$\mathcal{B} (\times 10^{-3})$
$\chi_{c0} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	2789 ± 66	9.30	$4.22 \pm 0.10 \pm 0.43$
$\chi_{c0} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	9031 ± 132	10.34	$8.61 \pm 0.13 \pm 0.94$
$\chi_{c0} \rightarrow \omega K^+ K^-$	1414 ± 42	8.04	$1.94 \pm 0.06 \pm 0.20$
$\chi_{c0} \rightarrow \phi \pi^+ \pi^- \pi^0$	538 ± 29	9.16	$1.18 \pm 0.07 \pm 0.13$
$\chi_{c1} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	5180 ± 75	10.21	$7.52 \pm 0.11 \pm 0.79$
$\chi_{c1} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	12256 ± 127	11.10	$11.46 \pm 0.12 \pm 1.29$
$\chi_{c1} \rightarrow \omega K^+ K^-$	628 ± 29	9.34	$0.78 \pm 0.04 \pm 0.08$
$\chi_{c1} \rightarrow \phi \pi^+ \pi^- \pi^0$	373 ± 26	10.50	$0.75 \pm 0.06 \pm 0.08$
$\chi_{c2} \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	4559 ± 71	9.76	$7.30 \pm 0.11 \pm 0.75$
$\chi_{c2} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	11189 ± 124	10.48	$11.69 \pm 0.13 \pm 1.31$
$\chi_{c2} \rightarrow \omega K^+ K^-$	512 ± 27	8.58	$0.73 \pm 0.04 \pm 0.08$
$\chi_{c2} \rightarrow \phi \pi^+ \pi^- \pi^0$	408 ± 28	9.88	$0.93 \pm 0.06 \pm 0.10$

TABLE VIII: Upper limits at the 90% C.L. on $\mathcal{B}(\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-)$ in the two η_c decay modes. N^{fit} is the number of events from the fits shown in Fig.3. In χ_{c2} case, N^{fit} includes the contribution from the peaking background $\psi(3686) \rightarrow \pi^+ \pi^- J/\psi, J/\psi \rightarrow \gamma \eta_c, \eta_c \rightarrow K_S^0 K^\pm \pi^\mp (K^+ K^- \pi^0)$.

Decay mode	N^{fit}	N^{up}	ϵ (%)	$\mathcal{B}^{\text{up}}(\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-)$ (%)	$\mathcal{B}^{\text{theory}}(\chi_{cJ} \rightarrow \eta_c \pi^+ \pi^-)$ (%)
$\chi_{c0} \rightarrow (K_S^0 K^\pm \pi^\mp) \pi^+ \pi^-$	0.0 ± 4.6	6.8	6.29	0.07	-
$\chi_{c0} \rightarrow (K^+ K^- \pi^0) \pi^+ \pi^-$	0 ± 15	33.6	6.82	0.41	-
$\chi_{c1} \rightarrow (K_S^0 K^\pm \pi^\mp) \pi^+ \pi^-$	18 ± 17	48.7	9.45	0.32	1.81 ± 0.26
$\chi_{c1} \rightarrow (K^+ K^- \pi^0) \pi^+ \pi^-$	6 ± 25	50.0	9.82	0.44	
$\chi_{c2} \rightarrow (K_S^0 K^\pm \pi^\mp) \pi^+ \pi^-$	77 ± 19	64.1	7.72	0.54	-
$\chi_{c2} \rightarrow (K^+ K^- \pi^0) \pi^+ \pi^-$	89 ± 26	105.4	7.83	1.23	-

Acknowledgments

The BESIII collaboration thanks the staff of BEPCII and the computing center for their hard efforts. This work is supported in part by the Ministry of Science and Technology of China under Contract No. 2009CB825200; National Natural Science Foundation of China (NSFC) under Contracts Nos. 10625524, 10821063, 10825524, 10835001, 10935007, 11125525; Joint Funds of the National Natural Science Foundation of China under Contracts Nos. 11079008, 11179007; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; CAS under Contracts Nos. KJCX2-YW-N29, KJCX2-YW-N45; 100 Talents Program of CAS; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; U. S. Department of Energy under Contracts Nos. DE-FG02-04ER41291, DE-FG02-91ER40682, DE-FG02-94ER40823; U.S.

National Science Foundation; University of Groningen (RuG); the Helmholtzzentrum fuer
Schwerionenforschung GmbH (GSI), Darmstadt; and WCU Program of National Research
Foundation of Korea under Contract No. R32-2008-000-10155-0.

-
- [1] N. Brambilla *et al.* (Quarkonium Working Group), CERN Yellow Report, CERN-2005-005.
 - [2] N. Brambilla *et al.* (Quarkonium Working Group), Eur. Phys. J. C **71**, 1534 (2011).
 - [3] T. M. Yan, Phys. Rev. D **22**, 1652 (1980).
 - [4] Y.-P. Kuang, S. F. Tuan, and T. M. Yan, Phys. Rev. D **37**, 1210 (1988).
 - [5] Y.-P. Kuang and T.-M. Yan, Phys. Rev. D **24**, 2874 (1981).
 - [6] Y.-P. Kuang, Front. Phys China **1**, 19 (2006).
 - [7] C. Cawfield *et al.* (CLEO Collaboration), Phys. Rev. D **73**, 012003 (2006).
 - [8] J. P. Lees *et al.* (BaBar Collaboration), arXiv:1206.2008 [hep-ex].
 - [9] Q. Liu and Y.-P. Kuang, Phys. Rev. D **75**, 054019 (2007). The predicted branching fraction is recalculated with the parameters in the model determined using the updated experimental data: $\Gamma(\psi(3686)) = (304 \pm 9) \text{ keV}$, $\mathcal{B}(\psi(3686) \rightarrow J/\psi\pi\pi) = (51.35 \pm 0.52)\%$, and $\mathcal{B}(\psi(3686) \rightarrow h_c\pi^0) \times \mathcal{B}(h_c \rightarrow \gamma\eta_c) = (4.35 \pm 0.57) \times 10^{-4}$. The total width of $\psi(3686)$ and the branching fraction of $\psi(3686) \rightarrow J/\psi\pi\pi$ are taken from PDG, while the $\mathcal{B}(\psi(3686) \rightarrow h_c\pi^0) \times \mathcal{B}(h_c \rightarrow \gamma\eta_c)$ is the combined result from the measurement in CLEO and BESIII Collaboration.
 - [10] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. D **81**, 052005 (2010).
 - [11] M. Ablikim *et al.* (BESIII Collaboration), Nucl. Instrum. Meth. A **614**, 345 (2010).
 - [12] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Meth. A **506**, 250 (2003).
 - [13] S. Jadach, B. F. L. Ward and Z. Was, Comp. Phys. Commu. **130**, 260 (2000); Phys. Rev. D **63**, 113009 (2001).
 - [14] <http://www.slac.stanford.edu/~lange/EvtGen/>; R. G. Ping *et al.*, Chinese Physics C **32**, 599 (2008).
 - [15] J. Beringer *et al.*, Phys. Rev. D **86**, 010001 (2012).
 - [16] J. C. Chen *et al.*, Phys. Rev. D **62**, 034003 (2000).
 - [17] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **108**, 222002 (2012).
 - [18] V. V. Anashin *et al.*, arXiv:1012.1694 [hep-ex].
 - [19] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **107**, 092001 (2011).
 - [20] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. Lett. **105**, 261801 (2010).
 - [21] M. Ablikim *et al.* (BESIII Collaboration), Phys. Rev. D **83**, 112005 (2011).
 - [22] R. E. Mitchell *et al.* (CLEO Collaboration), Phys. Rev. Lett. **102**, 011801 (2009).